



Lahar in a Jar

Instructions: Follow the procedures below to make a small lahar in a jar. You will mix known volumes of rock debris and water. Give your mixed lahar a run down a gutter or board provided by your teacher.


1. Place approximately 200 to 400 milliliters of "lahar" into a beaker. Break up any large clumps of dirt and debris. Record the exact volume here.

2. Make a prediction. How much water will be necessary to set the rock debris sample into motion as a small, in-the-jar lahar?

10 ml? 100 ml? Record your prediction.

of cookie dough and now resembles thick cake batter.

3. During this experiment, you will pour water into the beaker repeatedly, in increments of approximately 10 ml. In the space below, develop a procedure for keeping track of the amount of water that you tip during successive pours. Note: You may need to fill the graduated cylinder more than once during this experiment.







Lahar in a Jar-continued

5. Compare the total volume of water to the cumulative volume of lahar rocks and water. Use the space below to calculate the percent water required to form a lahar in the beaker. Record your result here.

ml water added Percent water = X 100 ml lahar deposit + ml of water added

- **6.** Determine whether the actual percent of water required to make a lahar is more or less than your prediction.
- 7. After completion of this experiment, preserve your sample for its run down a gutter or board as provided by your teacher. Explain why the slopes of Cascade volcanoes are an ideal location for the development of debris flows and lahars.

